

On limits of superstring in $AdS_5 \times S^5$

A.A. Tseytlin*

*Blackett Laboratory, Imperial College
London, SW7 2BZ, U.K.*

*Department of Physics, The Ohio State University
Columbus, OH 43210-1106, USA*

Abstract

The superstring action in $AdS_5 \times S^5$ depends on two parameters: the inverse string tension α' and the radius R . The “standard” AdS/CFT correspondence requires that the string coordinates are rescaled so that the action depends only on one combination of the two: $\sqrt{\lambda} = R^2/\alpha'$. Then $\lambda \rightarrow 0$ limit is equivalent to $R \rightarrow 0$ for fixed α' or to the zero-tension limit in $AdS_5 \times S^5$: $\alpha' \rightarrow \infty$ for fixed R . After reviewing previous work on the light cone superstring we explicitly obtain the $\lambda = 0$ form of its action. Its zero-mode part is the same as the superparticle action in $AdS_5 \times S^5$, and thus the $\lambda = 0$ string spectrum must include, as expected, the “protected” type IIB supergravity states. Following recent suggestions, it is conjectured that the spectrum of this tensionless string should as well contain higher spin massless states in AdS_5 . We also discuss the case of another parametrization of the string action which has straightforward $R \rightarrow \infty$ flat space limit but where $R \rightarrow 0$ and $\alpha' \rightarrow \infty$ limits are not equivalent. There $R \rightarrow 0$ corresponds to shrinking S^5 to zero the size and “freezing” the fluctuations of the radial coordinate of AdS_5 . This case is the basis of the “non-standard” AdS/CFT correspondence suggested in hep-th/0010106.

* Also at Lebedev Physics Institute, Moscow

1 Introduction

Further understanding of string theory – gauge theory duality in its simplest AdS/CFT case [1] depends on progress in solving string theory in $AdS_5 \times S^5$ with R-R flux. This string theory is described by a well-defined light-cone gauge Green-Schwarz action [2, 3], which, in contrast to its flat space counterpart [4], is non-linear. In particular, it contains complicated bosonic S^5 sigma model part as well as terms quartic in fermions. This complexity is not too surprising: the dual $\mathcal{N} = 4$ super Yang-Mills theory should be non-trivial even in the large N limit, with “observables” (dimensions of operators, OPE coefficients, entropy, etc.) being in general complicated functions of the ‘t Hooft coupling parameter $\lambda = g_{\text{YM}}^2 N$.^a

The light-cone $AdS_5 \times S^5$ string action is also different from the flat space GS action in that it does not have manifest 2-d Lorentz invariance. This is a generic feature of similar light-cone gauge string actions in curved space, and is not a problem of principle, in particular in the phase space approach, where the natural starting point is the light-cone Hamiltonian. One way to address the corresponding quantum problem is to expand the bosonic and fermionic string coordinates and momenta (which are functions of τ and periodic functions of σ) in Fourier modes in σ , getting a non-linear space-time supersymmetric quantum mechanical system with infinite number of bosonic and fermionic degrees of freedom. To analyse its spectrum one may first to restrict consideration to states that do not depend on S^5 directions and try to use numerical methods to deal with remaining non-linear terms in the Hamiltonian.

One may hope to simplify the problem of finding the string spectrum by considering some special limits of the parameters of the Hamiltonian – α' (inverse string tension) and R (radius of AdS_5 and S^5). In the general case of string in curved space of scale R , one may consider several independent limits when α' or R are sent separately to 0 or ∞ (under the assumption of a certain choice of momenta and coordinates which are held fixed in the limit) [3]. For example, in the particle theory limit ($\alpha' \rightarrow 0$), the string Hamiltonian reduces to the light-cone Hamiltonian for a superparticle in $AdS_5 \times S^5$ space [5], implying that the “massless” or zero-mode spectrum of the superstring coincides with the spectrum of type IIB supergravity compactified on S^5 [3]. Another is the null (tensionless) string limit $\alpha' \rightarrow \infty$, which, as in the flat space [6], is obtained by dropping terms in the Hamiltonian which contain derivatives with respect to the spatial world-sheet coordinate σ .

In the special context of the AdS/CFT correspondence, where the (large N) boundary conformal gauge theory should have spectrum of dimensions depending only on one parameter λ , the dual string theory should also effectively contain only one dynamical parameter – the effective tension proportional to $\sqrt{\lambda}$. To hope to establish the AdS/CFT correspondence one should thus make a special rescaling of the radial AdS string coordinate and its momentum so that the Hamiltonian will depend on α' and R only through one combination – $\lambda = R^4/\alpha'^2$.^b

^a Below we shall always consider the classical or first-quantized string theory, i.e. assume that $N \gg 1$.

^bNote that (combined also with different boundary conditions) this choice of coordinates will not allow one to directly recover the standard flat space 10-d type IIB superstring spectrum in the limit

The limit $\lambda \rightarrow 0$ has recently attracted some attention due to the conjecture that the free large N conformal $\mathcal{N} = 4$ SYM should be dual to a theory of massless higher spin fields in AdS_5 [7, 8, 9, 10, 11]. The basic observation is that the free gauge theory has conserved traceless conformal higher spin currents [12] that should be dual to (i.e. couple to the restriction to the boundary of) massless higher spins in AdS_5 . It would be important of course to see the presence of these massless higher spin states directly in the spectrum of the $AdS_5 \times S^5$ string in the limit $\lambda \rightarrow 0$.

With this motivation in mind, below we shall supplement the general expressions for the $AdS_5 \times S^5$ string action and Hamiltonian given in [3] with the explicit discussion of the $\lambda \rightarrow 0$ limit. For generality, we shall consider two different procedures of taking the $R \rightarrow 0$ limit, depending on how one scales the radial coordinate of the AdS space. The first one – relevant for the standard AdS/CFT correspondence – starts with the Hamiltonian depending on R and α' only through $\sqrt{\lambda} = R^2/\alpha'$. Here $R \rightarrow 0$ for fixed α' is equivalent to the zero tension limit $\alpha' \rightarrow \infty$ for fixed R , i.e. one effectively keeps the $AdS_5 \times S^5$ structure of space-time in the limit. Another $R \rightarrow 0$ limit can be taken in the action chosen in the form which has straightforward flat space limit when $R \rightarrow \infty$. Here $R \rightarrow 0$ means shrinking S^5 to zero size and also “freezing” the radial direction of AdS_5 . Starting with the string action written in covariant κ -symmetry gauge this second limit was considered previously in [13] in the context of an alternative version of the AdS/CFT ($AdS_5 \times S^5$ string – $\mathcal{N} = 4$ SYM) correspondence.

Before turning to the discussion of the string actions and their limits let us make few general comments on the “kinematics” of the AdS/CFT correspondence in the $\lambda = 0$ limit. The standard AdS/CFT correspondence in the sector of 5-d supergravity fields or boundary operators from the stress tensor supermultiplet can be described by coupling the $\mathcal{N} = 4$ SYM superconformal currents to $\mathcal{N} = 4$ conformal supergravity multiplet and then integrating over the $\mathcal{N} = 4$ SYM fields. The resulting large N effective action should then reproduce the 5-d gauged supergravity action (the quadratic and cubic terms in it, and, for large λ , also higher order terms) evaluated on the solution of the Dirichlet problem [1]. In particular, this was demonstrated explicitly at the quadratic level in the conformal graviton [14].^c

The same “induced action” procedure can be repeated for the free $\mathcal{N} = 4$ gauge theory which can be coupled to a higher spin generalization of the conformal supergravity multiplet. Coupling conserved traceless bilinear currents to the corresponding higher spin conformal fields, integrating out the free SYM fields (i.e. computing the logarithm of the determinants in the background) and expanding the resulting effective action to quadratic order in the background fields one may then compare this quadratic term to

$R \rightarrow \infty$, which of course is not a problem as it is not “seen” on the gauge theory side in any obvious limit.

^cCoupling the quantum SYM theory to background conformal supergravity multiplet and integrating out the SYM fields one finds the induced action for the conformal supergravity fields: $S = \int C_{mnkl} \ln(\epsilon^2 D^2) C_{mnkl} + \dots = c_1 \int (C_{mnkl}^2 + \dots) + \text{non-local terms}$. The quadratic and cubic terms in this action expanded in powers of the fields summarize information about protected correlators like $\langle T_{mn} T_{kl} \rangle$ and $\langle T_{mn} T_{kl} T_{sr} \rangle$ in conformal SYM theory. The same non-local action comes out of solving the Dirichlet problem in the 5-d $\mathcal{N} = 8$ gauged supergravity on the AdS_5 background.

the classical free action of the higher spin fields in AdS_5 evaluated on the solution of the Dirichlet problem. As in the case of the standard conformal supergravity, the agreement between the two actions is essentially kinematical, i.e. guaranteed by the symmetries. Related remarks and discussions of higher spin AdS/CFT correspondence appeared in [9, 10] and in [15, 16] (in light-cone gauge) and [11] (in covariant gauge).^d

More explicitly, the quadratic term in the local (logarithmically divergent) part of the corresponding 4-d effective action will be the free conformal higher spin action which has the following structure [18]

$$L = \sum_{s=1,2,\dots} \phi_s \partial^{2s} P_s \phi_s + \sum_{s=1/2,3/2,\dots} \psi_s \partial^{2s-1} \gamma^m \partial_m P_s \psi_s.$$

Here $s = 2$ and $s = 3/2$ terms correspond to the standard Weyl graviton and gravitino of conformal supergravity [19]. P_s are the transverse traceless projectors defined on totally symmetric fields [18]. Let us consider only the symmetric bosonic fields $\phi_s = (\phi_{m_1 \dots m_s})$.^e The fields ϕ_s have canonical dimension of $(\text{length})^{s-2}$ and can be coupled as $\int d^4x \phi_s(x) J_s(x)$ to the traceless currents $J_s = X \partial^s P_s X + \dots$ which are conserved when X is on shell (X with canonical dimension $(\text{length})^{-1}$ are the scalars of the $\mathcal{N} = 4$ vector multiplet). Starting with $L = X \partial^2 X + \phi_s(x) X \partial^s P_s X + \dots$ and integrating out the SYM fields gives the induced action $\Gamma[\phi_s] \sim \ln \det(-\partial^2 + \phi_s \partial^s P_s)$. The quadratic term in this action will be proportional to $\int d^4p \phi_s(p) p^{2s} \ln(\epsilon^2 p^2) P_s \phi_s(-p)$ or, in coordinate representation, $\int \phi_s(x) \partial^{2s} \ln(\epsilon^2 \partial^2) P_s \phi_s$, with the local (divergent) part being the above pure conformal higher spin action. This quadratic term in $\Gamma[\phi_s]$ may be written also as $\int d^4x d^4x' \frac{\phi_s(x) \hat{P}_s(x-x') \phi_s(x')}{(|x-x'|^2 + \epsilon^2)^{s+2}}$. On symmetry grounds, the same non-local functional must follow from the solution of the Dirichlet problem for the corresponding massless higher spin in AdS_5 .^f

2 $R \rightarrow 0$ limit in string action in covariant gauge

We shall start with a heuristic discussion of the $R \rightarrow 0$ limit taken directly in the $AdS_5 \times S^5$ superstring action in covariant gauge.

The superstring Lagrangian [23] may be written in “4+6” parametrization of $AdS_5 \times$

^dNote that the $\lambda = 0$ theory will contain several currents or conformal fields of the same spin. The corresponding generalization of conformal supergravity should thus include, in particular, several [7] spin 2 conformal gravitons. Such theories were recently discussed in [17].

^eInteracting higher spin conformal field theories should exist in flat space and have dimensionless cubic, etc., couplings [20, 21]. Containing higher derivatives in the kinetic terms they non-unitary (like conformal supergravity or Weyl gravity), but non-unitarity is not an issue in the present context. Here such theory is viewed as an induced (non-local) effective field theory which is simply a way of encoding the information about the correlators of higher spin conformal currents of the SYM theory. Refs. [20, 21] speculated about possible connections between conformal higher spins in flat space, massless higher spins in AdS, and string theory, but they were not anticipating the $4 \rightarrow 5$ holographic jump in the dimension essential for the AdS/CFT correspondence.

^fIncluding non-linear in ϕ_s terms in the coupling to SYM fields will not change the value of the conformal current correlators at separated points but is necessary for maintaining manifest higher spin symmetries in the induced action. Let us note in passing that viewing the $\mathcal{N} = 4$ conformal supergravity action as the local part of the induced action generated by integrating out the $\mathcal{N} = 4$ SYM fields in the conformal supergravity background implies that the conformal supergravity action must contain the Weyl tensor term C_{mnkl}^2 without any extra scalar-field dependent prefactor, cf. [22, 18].

S^5 corresponding to the metric ($a = 0, \dots, 3$, $M = 1, \dots, 6$, $Y^M = e^\phi u^M$, $u^M u^M = 1$)

$$ds^2 = Y^2 dx^a dx^a + R^2 Y^{-2} dY^M dY^M = e^{2\phi} dx^a dx^a + R^2 d\phi^2 + R^2 du^M du^M . \quad (2.1)$$

This metric has the standard flat space limit when $R \rightarrow \infty$ (setting $\phi = \varphi/R$ but without need to rescale x^a). Another parametrization is based on rescaling Y by R (i.e. shifting ϕ by $\ln R$). Then R^2 is the factor in front of the whole 10-d metric

$$ds^2 = R^2 (Y^2 dx^a dx^a + Y^{-2} dY^M dY^M) = R^2 (e^{2\phi} dx^a dx^a + d\phi^2 + du^M du^M) . \quad (2.2)$$

This second choice is the basis for the standard AdS/CFT correspondence where the string action will then depend on R through $\sqrt{\lambda} = R^2/\alpha'$. This choice means that all distances, including the world-sheet ones are effectively measured in terms of R .

Interpreting the $AdS_5 \times S^5$ isometry supergroup $PSU(2, 2|4)$ as the $\mathcal{N} = 4$ superconformal group in 4 dimensions, it is natural to split the fermionic generators into 4 standard supergenerators Q_i and 4 special conformal supergenerators S_i (we suppress the 4-d spinor indices). The associated superstring coordinates will be denoted as θ_i and η_i [2, 3]. The 4-d Lorentz covariant κ -symmetry “S-gauge” of [2] corresponds to setting all η_i fields to zero. The resulting Lagrangian has the following simple structure [2]⁹

$$L = -\frac{1}{2} \sqrt{g} [Y^2 (\partial_\mu x^a - i\theta \sigma^a \partial_\mu \theta)^2 + R^2 Y^{-2} \partial^\mu Y^M \partial_\mu Y^M] - i\epsilon^{\mu\nu} R \partial_\mu Y^M \theta \rho^M \partial_\nu \theta . \quad (2.3)$$

Here

$$i\theta \sigma^a \partial_\mu \theta \equiv i\theta_{\dot{a}i} \sigma^{a\dot{a}b} \partial_\mu \theta_b^i + h.c. , \quad i\theta \rho^M \partial_\nu \theta \equiv i\theta^{\dot{a}i} \rho_{ij}^M \partial_\nu \theta_{\dot{a}}^j + h.c. .$$

We have written the Lagrangian corresponding to the metric (2.1), where Y^M is dimensionless (x^a and R have dimension of length and the fermionic coordinates θ have dimension (length)^{1/2}). The resulting string action $I = \frac{1}{2\pi\alpha'} \int d^2\sigma L$ is then dimensionless. To see that the action (2.3) has the correct flat space limit one is to set $\phi = \varphi/R$ and send R to infinity.

The action corresponding to (2.2) is obtained by $Y^M \rightarrow RY^M$:

$$L = R^2 \left\{ -\frac{1}{2} \sqrt{g} [Y^2 (\partial_\mu x^a - i\theta \sigma^a \partial_\mu \theta)^2 + Y^{-2} \partial^\mu Y^M \partial_\mu Y^M] - i\epsilon^{\mu\nu} \partial_\mu Y^M \theta \rho^M \partial_\nu \theta \right\} . \quad (2.4)$$

As discussed in the Introduction, the way to take the $R \rightarrow 0$ limit is in general not unique: the result depends on additional assumptions about which coordinates and arguments (momenta) of external states are held fixed in the limit. For example, starting with (2.3) or (2.4) gives apparently different results. The action corresponding to (2.4) depends on R through the dimensionless ratio $\lambda^{1/4} = R/\sqrt{\alpha'}$, so that $R \rightarrow 0$ corresponds to $\lambda \rightarrow 0$

⁹The actions in [25, 26, 27] have equivalent forms, corresponding to particular choices of the 10-d Dirac matrix representation. We use the following notation: the 4-d indices are $a, b = 0, 1, 2, 3$; $SO(6)$ indices are $M, N = 1, \dots, 6$; $SU(4)$ indices are $i, j = 1, 2, 3, 4$; the 2-d indices are $\mu, \nu = 0$. We use “chiral” representations for the 4-d and 6-d Dirac matrices, $\gamma^a = \begin{pmatrix} 0 & \sigma^a \\ \bar{\sigma}^a & 0 \end{pmatrix}$, $\gamma^M = \begin{pmatrix} 0 & \rho^{Mij} \\ \rho_{ij}^M & 0 \end{pmatrix}$, with $(\rho^M)^{ij} \equiv -(\rho_{ij}^M)^*$. $\mathbf{a}, \mathbf{b} = 1, 2$ are the $sl(2, C)$ (i.e. 4-d spinor) indices and $\theta_{\mathbf{a}i}^\dagger = -\theta_{\mathbf{a}}^i$, $\theta_i^{\mathbf{a}\dagger} = \theta^{\mathbf{a}i}$. The 10-d spinors are split as (θ^i, θ_i) .

and thus is also equivalent to $\alpha' \rightarrow \infty$ with R (and external momenta) kept finite in this limit.

In the rest of this section we shall follow [13] and discuss taking the $R \rightarrow 0$ limit in the case of (2.3). The direct result of setting $R = 0$ in (2.3) is

$$L_{R \rightarrow 0} = -\frac{1}{2}\sqrt{g} Y^2 (\partial_\mu x^a - i\theta\sigma^a\partial_\mu\theta)^2, \quad (2.5)$$

suggesting that six Y^M coordinates effectively “decouple”. This truncated action still has the expected $\mathcal{N} = 4$ global linear supersymmetry and 4-d conformal invariance (Y is still to be integrated over).

The action (2.5) does not of course define a consistent critical string theory and should be supplemented with a more systematic procedure of taking the $R \rightarrow 0$ limit in the path integral with external operator insertions. Since the coefficient of the kinetic term of Y^M directions goes to zero in the limit, that means that if the external operator depends on non-constant modes of Y^M then the $R \rightarrow 0$ limit will be singular. To be able to define this limit, one should thus restrict consideration to external states that do not depend on non-constant modes in radial AdS_5 and all S^5 directions. However, one should still integrate over the non-constant modes of Y^M in the path integral, and this is essential to ensure that the resulting string theory is consistent. Indeed, the $AdS_5 \times S^5$ string sigma model (2.3) is conformal and has the right central charge for any R [23, 24], so taking the limit after doing the integral over all of the fields including the non-constant modes of Y^M should not affect these properties.

The constant modes of Y^M should in general couple to external states and the integral over them will be non-trivial, i.e. we should get, symbolically,^{*h*}

$$\langle F \rangle = \int d^6 Y_0 M(Y_0) \int Dx D\theta e^{iI[x,\theta,Y_0]} F[x,\theta,Y_0], \quad (2.6)$$

where Y_0 is the constant part of Y . We have added the measure factor $M(Y_0) = \sqrt{G_{10}} = Y^{-2}$ (but did not explicitly split x and θ on constant and non-constant modes). Assuming further that external states do not depend on S^5 angles (i.e. F is $SO(6)$ symmetric), we get (setting $Y_0 = e^{\phi_0}$, $T = \frac{1}{2\pi\alpha'}$)

$$\langle F \rangle \sim \int d\phi_0 e^{4\phi_0} \int Dx D\theta e^{-\frac{i}{2}T \int d^2\sigma \sqrt{g} e^{2\phi_0} (\partial_\mu x^a - i\theta\sigma^a\partial_\mu\theta)^2} F[x,\theta,\phi_0]. \quad (2.7)$$

The integral over the constant ϕ_0 should then ensure the target space scale invariance in 4 dimensions, i.e. the symmetry under rescalings of x^a .

To summarize, in the particular $R \rightarrow 0$ limit in (2.3) corresponding to (2.1) the string is effectively allowed to fluctuate only in 4 space-time dimensions (as well as in fermionic dimensions), and the integration over the constant mode ϕ_0 of the radial AdS_5 direction produces averaging over the effective string tension ($\mathcal{T} \equiv T e^{2\phi_0}$)

$$\langle F \rangle \sim \int_0^\infty d\mathcal{T} \mathcal{T}^{3/2} \int Dx D\theta e^{-\frac{i}{2}\mathcal{T} \int d^2\sigma \sqrt{g} (\partial_\mu x^a - i\theta\sigma^a\partial_\mu\theta)^2} F[x,\theta,\mathcal{T}]. \quad (2.8)$$

^{*h*}Again, it is understood that one has first integrated out all non-zero modes of Y^M to cancel the conformal anomaly and ensure that the model is finite.

If instead one starts with the action (2.4) then the $R \rightarrow 0$ limit is equivalent to the zero tension limit $T \rightarrow 0$ and is apparently a strong-coupling limit in all 10 directions (see also [28] for related remarks). In general, the covariant gauge action is not the best starting point for the investigation of the $R \rightarrow 0$ limit because it does not allow one to find the spectrum of “small” strings (the kinetic term of fermions is not well defined). We shall therefore turn now to the discussion of the $R \rightarrow 0$ limit in the light-cone gauge.

3 Superstring action in fermionic light-cone gauge

In flat space, the superstring light-cone gauge fixing includes the fermionic light-cone gauge choice (i.e. fixing the κ -symmetry by the $\gamma^+ \theta^I = 0$ condition), and the bosonic light-cone gauge choice (i.e. using the conformal gaugeⁱ $\sqrt{g}g^{\mu\nu} = \eta^{\mu\nu}$ and fixing the residual conformal diffeomorphism symmetry by $x^+(\tau, \sigma) = p^+ \tau$). Fixing the fermionic light-cone gauge produces already a substantial simplification of the flat-space GS action: it becomes quadratic in θ . Similarly, in $AdS_5 \times S^5$ one is able to find a light-cone κ -symmetry gauge [2] in which the fermion kinetic term has the structure $\partial x^+ \theta \partial \theta$. It thus involves only one combination $-x^+$ of the 4-d coordinates, so that the non-degeneracy of the fermion kinetic term does not depend on a choice of specific string background in transverse directions. To simplify the fermion kinetic term further one may choose next the bosonic light-cone gauge $x^+ = p^+ \tau$. Fixing the bosonic part of the light-cone gauge and derivation of the resulting light-cone Hamiltonian was described in detail in [3] (see also [30]) and will be discussed in the next section.

Let us first review the form of the string action found after fixing the light-cone κ -symmetry gauge. In contrast to the Lorentz covariant “S-gauge” where 16 fermions η_i are set equal to zero, the light-cone gauge used in [2] corresponds to setting to zero “half” of the 16 θ_i and “half” of the 16 η_i fermionic coordinates (“half” is defined with respect to $SO(1, 1)$ rotations in the light-cone directions). The remaining fermions (which will be again denoted by θ_i, η_i) from now on will be simply 4+4 complex anticommuting variables carrying no extra Lorentz spinor indices. For comparison, the flat space GS action in the light-cone gauge ($\gamma^+ \theta = 0$) [4], written in a similar parametrization of the 16 fermionic coordinates, has the following structure (suppressing the $SU(4)$ indices)

$$L = -\frac{1}{2}\sqrt{g}(\partial_\mu x)^2 - \left[\frac{i}{2}\sqrt{g}\partial^\mu x^+(\theta\partial_\mu\theta + \eta\partial_\mu\eta) - \epsilon^{\mu\nu}\partial_\mu x^+\eta\partial_\nu\theta + h.c. \right]. \quad (3.9)$$

This form of the original GS Lagrangian is indeed the flat space ($R \rightarrow \infty$) limit of the light-cone $AdS_5 \times S^5$ Lagrangian of [2] written in the parametrization used in (2.1) (cf. (2.3))

$$L = -\sqrt{g}\left[Y^2(\partial^\mu x^+\partial_\mu x^- + \partial^\mu x\partial_\mu \bar{x}) + \frac{1}{2}R^2Y^{-2}(\partial_\mu Y^M + iR^{-2}\eta\rho^{MN}\eta Y^N Y^2\partial_\mu x^+)^2\right] \\ - \left(\frac{i}{2}\sqrt{g}g^{\mu\nu}Y^2\partial_\mu x^+[\theta^i\partial_\nu\theta_i + \eta^i\partial_\nu\eta_i + \frac{i}{2}R^{-2}Y^2\partial_\nu x^+(\eta^2)^2]\right)$$

ⁱBelow we shall use the Minkowski signature 2-d world sheet metric $g_{\mu\nu}$ with $g \equiv -\det g_{\mu\nu}$.

$$- \epsilon^{\mu\nu} |Y| \partial_\mu x^+ \eta^i \rho_{ij}^M Y^M (\partial_\nu \theta^j - i R^{-1} \sqrt{2} |Y| \eta^j \partial_\nu x) + h.c.) . \quad (3.10)$$

x^a are decomposed into the light-cone (x^+, x^-) and 2 complex transverse coordinates $x_\perp = (x, \bar{x})$, i.e. $x^\pm = \frac{1}{\sqrt{2}}(x^3 \pm x^0)$; $x, \bar{x} = \frac{1}{\sqrt{2}}(x^1 \pm ix^2)$.^j The superconformal algebra $psu(2, 2|4)$ implies that the 16 physical Grassmann variables – “linear” θ^i and “nonlinear” η^i and their hermitian conjugates θ_i and η_i which transform in the fundamental representations of $SU(4)$ are related to the Poincaré and the conformal supersymmetry in the light-cone gauge description of the boundary theory.^k

The dependence on the radius R is consistent with θ and η having dimensions (length)^{1/2} and Y^M being dimensionless in (2.1). The η -fermionic terms with R^{-2} factors correspond to the connection (and R-R coupling) and curvature terms that go away in the flat space limit ($|Y| \rightarrow 1$) where one recovers (3.9).^l

Written in terms of the radial direction ϕ of AdS_5 and unit 6-d vector u^M parametrizing S^5 the Lagrangian (3.10) becomes ($Y^M = e^\phi u^M$, $|Y| = e^\phi$) [2]

$$\begin{aligned} L = & -\sqrt{g} e^{2\phi} (\partial^\mu x^+ \partial_\mu x^- + \partial^\mu x \partial_\mu \bar{x}) \\ & - \frac{1}{2} \sqrt{g} R^2 [\partial^\mu \phi \partial_\mu \phi + (\partial_\mu u^M + i R^{-2} \eta \rho^{MN} \eta u^N e^{2\phi} \partial_\mu x^+)^2] \\ & - \left(\frac{i}{2} \sqrt{g} g^{\mu\nu} e^{2\phi} \partial_\mu x^+ [\theta^i \partial_\nu \theta_i + \eta^i \partial_\nu \eta_i + \frac{i}{2} R^{-2} e^{2\phi} \partial_\nu x^+ (\eta^2)^2] \right. \\ & \left. - \epsilon^{\mu\nu} e^{2\phi} \partial_\mu x^+ \eta^i \rho_{ij}^M u^M (\partial_\nu \theta^j - i R^{-1} \sqrt{2} e^\phi \eta^j \partial_\nu x) + h.c. \right) . \end{aligned} \quad (3.11)$$

Starting with the action (3.11), to get a regular $R \rightarrow 0$ limit we need to deal with blowing up of the fermionic terms with R^{-1} factors. One option is to demand that external states have no dependence on η , i.e. to effectively freeze η to zero.^m By formally setting $R = 0$ in the action we are then left with

$$L_{R \rightarrow 0} = -\sqrt{g} e^{2\phi} \left[\partial^\mu x^+ \partial_\mu x^- + \partial^\mu x \partial_\mu \bar{x} + \partial^\mu x^+ \left(\frac{i}{2} \theta^i \partial_\mu \theta_i + h.c. \right) \right] . \quad (3.12)$$

This is the light-cone gauge analog of the “naive” action (2.5). As in the previous section, to get a well-defined limit we should again average only quantities that do not have dependence on non-constant modes of ϕ , u^M (which should be integrated out before taking $R \rightarrow 0$). Since the integral over ϕ will be effectively restricted to constant values it is not problem to fix the standard bosonic light-cone gauge $x^+ = p^+ \tau$. Note that in contrast to the standard flat space GS action the resulting fermionic term in the action

^jThe matrices ρ^M (off-diagonal blocks of 6-d Dirac matrices in chiral representation) satisfy: $\rho_{ij}^M = -\rho_{ji}^M$, $(\rho^M)^{il} \rho_{lj}^N + (\rho^N)^{il} \rho_{lj}^M = 2\delta^{MN} \delta_j^i$, $(\rho^M)^{ij} \equiv -(\rho_{ij}^M)^*$, and $\rho^{MN} \equiv \rho^{[M} \rho^{\dagger N]}$, i.e. $\rho^{MNi} = \frac{1}{2} (\rho^M)^{il} \rho_{lj}^N - (M \leftrightarrow N)$. Also, $\theta_i^\dagger = \theta^i$, $\eta_i^\dagger = \eta^i$, $\theta^2 \equiv \theta^i \theta_i$, $\eta^2 \equiv \eta^i \eta_i$.

^kThe action is invariant under shifting $\theta \rightarrow \theta + \epsilon$ (supplemented with a shift of x^-).

^lThe presence of the η^4 term reflects the curvature of the background; the ‘extra’ $O(\eta^2)$ terms have the interpretation of the couplings to the R-R 5-form background [23].

^mA more systematic procedure is to introduce $\zeta \equiv R^{-1} \eta$ which will be fixed in the $R \rightarrow 0$ limit. Then the ζ -dependence decouples in the $R \rightarrow 0$ limit, i.e. this limit is regular provided the external states do not depend on ζ .

will miss the spatial derivative part, i.e. will not be 2-d Lorentz covariant. As discussed in [2, 3], this is a general feature of the light-cone gauge actions in curved AdS-type spaces.

Let us now turn to the case of the action which is relevant for the standard AdS/CFT correspondence, i.e. the one associated the metric (2.2) and thus related to (3.11) by $e^\phi \rightarrow Re^\phi$,

$$L = R^2 \left\{ -\sqrt{g}e^{2\phi}(\partial^\mu x^+ \partial_\mu x^- + \partial^\mu x \partial_\mu \bar{x}) - \frac{1}{2}\sqrt{g}[\partial^\mu \phi \partial_\mu \phi + (\partial_\mu u^M + i\eta \rho^{MN} \eta u^N e^{2\phi} \partial_\mu x^+)^2] \right. \\ \left. - \left(\frac{i}{2}\sqrt{g}g^{\mu\nu} e^{2\phi} \partial_\mu x^+ [\theta^i \partial_\nu \theta_i + \eta^i \partial_\nu \eta_i + \frac{i}{2}e^{2\phi} \partial_\nu x^+ (\eta^2)^2] \right. \right. \\ \left. \left. - \epsilon^{\mu\nu} e^{2\phi} \partial_\mu x^+ \eta^i \rho_{ij}^M u^M (\partial_\nu \theta^j - i\sqrt{2}e^\phi \eta^j \partial_\nu x) + h.c. \right) \right\}. \quad (3.13)$$

As in the covariant gauge action (2.4) of the previous section, here $R \rightarrow 0$ is the strong-coupling limit in all directions, and there seems to be no natural simplification of the action. In such cases it is more appropriate to use the (light-cone gauge) phase space formulation to the discussion of which we now turn.

4 Bosonic light-cone gauge fixing in AdS space

To eliminate the ∂x^+ -factors from the fermion kinetic terms in (3.11) or (3.13) one is to fix the bosonic light-cone gauge. In flat space this may be done by fixing the conformal gauge $\sqrt{g}g^{\mu\nu} = \eta^{\mu\nu}$, and then noting that one can fix the residual conformal diffeomorphism symmetry on the plane by choosing $x^+(\tau, \sigma) = p^+ \tau$. In the case of the AdS-type curved spaces, the bosonic light-cone gauge $x^+ = p^+ \tau$ can not be combined with the standard conformal gauge $\sqrt{g}g^{\mu\nu} = \eta^{\mu\nu}$. One needs instead to impose a condition on $g_{\mu\nu}$ that breaks the manifest 2-d Lorentz symmetry. One consistent gauge choice is [2]: $e^{2\phi} \sqrt{g}g^{00} = -1$, $x^+ = p^+ \tau$. Using these conditions in the action (3.11) and integrating over x^- and the remaining components of the 2-d metric one finds the explicit form of the action which is equivalent to the one obtained in [3] in the phase space approach. A closely related alternative originally suggested in [29] is a modification of the conformal gauge $\sqrt{g}g^{ab} = \text{diag}(-e^{-2\phi}, e^{2\phi})$ which, as in flat space, it turns out to be possible to supplement with the third light-cone gauge condition $x^+ = p^+ \tau$.ⁿ

The resulting action has AdS_5 radial direction e^ϕ factors coupled differently to ∂_0 and ∂_1 derivative terms, and the S^5 part of the action is also coupled to ϕ . In the absence of manifest 2-d Lorentz symmetry it is natural to use the phase space formulation of the light-cone gauge fixed theory. The coordinate space Polyakov approach based on any of the above gauges is equivalent to the phase space GGRT approach based on fixing the diffeomorphisms by $x^+ = \tau$, $P^+ = p^+ = \text{const}$ [3].

To illustrate the derivation of the phase space Lagrangian let us first ignore the S^5 and fermionic parts in (3.13), i.e. start with

$$\mathcal{L} \equiv TL = -Th^{\mu\nu}(\partial_\mu x^+ \partial_\nu x^- + \partial_\mu x \partial_\nu \bar{x} + \frac{1}{2}e^{-2\phi} \partial_\mu \phi \partial_\nu \phi), \quad (4.14)$$

ⁿEquivalent action is found also using the gauge [31] $g_{01} = 0$, $x^+ = p^+ \tau$.

$$T \equiv TR^2 = \frac{R^2}{2\pi\alpha'} = \frac{\sqrt{\lambda}}{2\pi}, \quad h^{\mu\nu} \equiv \sqrt{g}g^{\mu\nu}e^{2\phi}. \quad (4.15)$$

Introducing the momenta $\mathcal{P}^a = T \frac{\partial L}{\partial \dot{x}^a}$ for the light-cone coordinates (x^+, x^-) and the two transverse coordinates $x_\perp = (x, \bar{x})$ and the momentum Π for the radial direction ϕ we get

$$\begin{aligned} \mathcal{L} = & \dot{x}_\perp \mathcal{P}_\perp + \dot{\phi} \Pi + \dot{x}^+ \mathcal{P}^- + \dot{x}^- \mathcal{P}^+ + \frac{1}{2h^{00}} \left[T^{-1} \mathcal{P}_\perp^2 + T^{-1} \mathcal{P}^+ \mathcal{P}^- \right. \\ & \left. + T e^{4\phi} (\dot{x}_\perp^2 + 2\dot{x}^+ \dot{x}^-) + e^{2\phi} (T^{-1} \Pi^2 + T \dot{\phi}^2) \right] + \frac{h^{01}}{h^{00}} (\dot{x}_\perp \mathcal{P}_\perp + \dot{\phi} \Pi + \dot{x}^+ \mathcal{P}^- + \dot{x}^- \mathcal{P}^+), \end{aligned} \quad (4.16)$$

where $1/h^{00}$ and h^{01}/h^{00} play the role of the Lagrange multipliers imposing two constraints. Choosing the light-cone gauge $x^+ = \tau$, $\mathcal{P}^+ = p^+ = \text{const}$ and integrating over $\mathcal{P}^-$ we get the relation $h^{00} = -T^{-1}p^+$. The expression for $\mathcal{P}^-$ follows from the $1/h^{00}$ constraint after using the h^{01}/h^{00} constraint. The resulting light-cone gauge Hamiltonian density $\mathcal{H}$ is [3]

$$\mathcal{H} = -\mathcal{P}^- = \frac{1}{2p^+} \left[\mathcal{P}_\perp^2 + e^{4\phi} T^2 \dot{x}_\perp^2 + e^{2\phi} (\Pi^2 + T^2 \dot{\phi}^2) \right]. \quad (4.17)$$

The constraint determining x^- is $p^+ \dot{x}^- + \mathcal{P}_\perp \dot{x}_\perp + \dot{\phi} \Pi = 0$. The coordinates and momenta satisfy the usual Poisson bracket relation $\{x(\tau, \sigma), \mathcal{P}(\tau, \sigma')\} = \delta(\sigma - \sigma')$. Integrating over the momenta $\mathcal{P}_\perp$ and Π , one finds the corresponding light-cone gauge Lagrangian

$$L = \frac{p^+}{2} \left(\dot{x}_\perp^2 - e^{4\phi} T^2 (p^+)^{-2} \dot{x}_\perp^2 + e^{-2\phi} [\dot{\phi}^2 - e^{4\phi} T^2 (p^+)^{-2} \phi'^2] \right). \quad (4.18)$$

As usual, the explicit dependence on p^+ can be absorbed into redefinition $\sigma \rightarrow \sigma/p^+$ in the action. The analog of (4.17) in the case of (2.1), i.e. the action (3.11), is obtained by $e^\phi \rightarrow R^{-1}e^\phi$,

$$\mathcal{H} = -\mathcal{P}^- = \frac{1}{2p^+} \left[\mathcal{P}_\perp^2 + e^{4\phi} T^2 \dot{x}_\perp^2 + R^{-2} e^{2\phi} (\Pi^2 + T^2 R^4 \dot{\phi}^2) \right]. \quad (4.19)$$

In this case $\mathcal{H}$ depends separately on the tension T and the radius R , and so one may consider several limits (assuming other arguments of $\mathcal{H}$ are kept fixed): (i) $T \rightarrow \infty$, $R = \text{fixed}$: particle in AdS; (ii) $T \rightarrow 0$, $R = \text{fixed}$: “standard” tensionless string in AdS; (iii) $R \rightarrow \infty$, $T = \text{fixed}$: flat space; (iv) $R \rightarrow 0$, $T = \text{fixed}$: string in “zero radius” AdS. To get finite result in the particle limit $T \rightarrow \infty$ one must set $x' = 0$, $\phi' = 0$, i.e. assume that the string shrinks to a point, $H_{T \rightarrow \infty} = \frac{1}{2}(\mathcal{P}_\perp^2 + e^{2\phi} \Pi^2)$. The limit $R \rightarrow \infty$ in (4.19) is defined by setting $\phi = \varphi/R$, $\Pi = Rp$; this preserves the Poisson bracket and gives the standard regular flat space expression. To define the limit $R \rightarrow 0$ in (4.19) we must (as in the previous section) restrict consideration to states that do not depend on Π , i.e. have zero momentum in the radial AdS direction, $\mathcal{H}_{R \rightarrow 0} = \frac{1}{2p^+} (\mathcal{P}_\perp^2 + T^2 e^{4\phi} \dot{x}_\perp^2)$.

In the case of (4.17) the flat space limit is not reached directly, and there are only two independent limits: $T \rightarrow 0$ and $T \rightarrow \infty$. Here the limit $R \rightarrow 0$ for fixed T is equivalent to $T \sim \sqrt{\lambda} \rightarrow 0$ and thus to $T \rightarrow 0$ for fixed R . It thus corresponds (as the usual zero-tension limit in flat space) to dropping σ -derivative terms in the action and Hamiltonian, i.e.

$$\mathcal{H}_{\lambda \rightarrow 0} = \frac{1}{2p^+} (\mathcal{P}_\perp^2 + e^{2\phi} \Pi^2). \quad (4.20)$$

5 Superstring light cone action and its $\lambda \rightarrow 0$ limit

Let us now add the dependence on S^5 and fermionic directions. Repeating the above bosonic light-cone gauge fixing procedure for the superstring action in (3.13) we get [3]

$$\begin{aligned} \mathcal{L} = & \mathcal{P}_\perp \dot{x}_\perp + \Pi \dot{\phi} + \mathcal{P}_M \dot{u}^M + \frac{i}{2} p^+ (\theta^i \dot{\theta}_i + \eta^i \dot{\eta}_i - h.c.) - \mathcal{H} \\ & - T(p^+)^{-1} h^{01} \left[p^+ \dot{x}^- + \mathcal{P}_\perp \dot{x}_\perp + \Pi \dot{\phi} + \mathcal{P}_M \dot{u}^M + \frac{i}{2} p^+ (\theta^i \dot{\theta}_i + \eta^i \dot{\eta}_i - h.c.) \right], \end{aligned} \quad (5.21)$$

where the light-cone Hamiltonian density is

$$\begin{aligned} \mathcal{H} = -\mathcal{P}^- = & \frac{1}{2p^+} \left(\mathcal{P}_\perp^2 + T^2 e^{4\phi} \dot{x}_\perp^2 + e^{2\phi} [\Pi^2 + T^2 \dot{\phi}^2 + \mathcal{P}^M \mathcal{P}^M + T^2 \dot{u}^M \dot{u}^M \right. \\ & \left. + p^{+2} (\eta^2)^2 + 2ip^+ \eta_i (\rho^{MN})^i_j \eta^j u^M \mathcal{P}^N \right] - T e^{2\phi} u^M [\eta^i \rho_{ij}^M (\dot{\theta}^j - i\sqrt{2} e^\phi \eta^j \dot{x}) + h.c.] . \end{aligned} \quad (5.22)$$

Here the odd part of the phase space is represented by θ^i, η^i considered as fermionic coordinates and θ_i, η_i considered as fermionic momenta. h^{01} imposes the reparametrization constraint, and $\mathcal{P}^M$ is the momentum corresponding to the unit 6-vector u^M , $u^M \mathcal{P}^M = 0$.

In the particle theory limit, i.e. $T \rightarrow \infty$, implying that all spatial derivatives are to be set to zero, (5.22) reduces [3] to the (classical limit of) quantum light-cone Hamiltonian for a superparticle in $AdS_5 \times S^5$ space [5].^o

The equivalent expressions for the string Lagrangian and Hamiltonian written in terms of 6 Cartesian coordinates Y^M and the associated (now unconstrained) momenta $\mathcal{P}_M$, i.e. corresponding to the Lagrangian (3.10) but with the “standard AdS/CFT” rescaling $Y^M \rightarrow RY^M$, $\mathcal{P}_M \rightarrow R^{-1}\mathcal{P}_M$ are

$$\mathcal{L} = \mathcal{P}_\perp \dot{x}_\perp + \mathcal{P}_M \dot{Y}^M + \frac{i}{2} p^+ (\theta^i \dot{\theta}_i + \eta^i \dot{\eta}_i - h.c.) - \mathcal{H}, \quad (5.23)$$

$$\begin{aligned} \mathcal{H} = & \frac{1}{2p^+} \left(\mathcal{P}_\perp^2 + T^2 Y^4 \dot{x}_\perp^2 + Y^4 \mathcal{P}_M \mathcal{P}_M + T^2 \dot{Y}^M \dot{Y}^M \right. \\ & \left. + Y^2 [p^{+2} (\eta^2)^2 + 2ip^+ \eta \rho^{MN} \eta Y_M \mathcal{P}_N] \right) \\ & - T |Y| Y^M [\eta \rho^M (\dot{\theta} - i\sqrt{2} |Y| \eta \dot{x}) + h.c.] . \end{aligned} \quad (5.24)$$

Here also $p^+ \dot{x}^- + \mathcal{P}_\perp \dot{x}_\perp + \mathcal{P}_M \dot{Y}^M + \frac{i}{2} p^+ (\theta^i \dot{\theta}_i + \eta^i \dot{\eta}_i - h.c.) = 0$.

The $\lambda \rightarrow 0$ limit in (5.24) is the same as $T \rightarrow 0$, i.e. we get

$$\mathcal{H}_{\lambda \rightarrow 0} = \frac{1}{2p^+} \left(\mathcal{P}_\perp^2 + Y^4 \mathcal{P}_M \mathcal{P}_M + Y^2 [p^{+2} (\eta^2)^2 + 2ip^+ \eta \rho^{MN} \eta Y_M \mathcal{P}_N] \right). \quad (5.25)$$

In contrast to the tensionless string limit in flat space where one gets an infinite set of “decoupled” modes, here the S^5 and fermionic terms introduce non-trivial interactions

^oAs a result, the “massless” (zero-mode) spectrum of the superstring coincides indeed with the spectrum of type IIB supergravity compactified on S^5 . The vertex operators for the supergravity states are then obtained by solving the linearized supergravity equations (or 1-st quantized superparticle state equations) in $AdS_5 \times S^5$ background.

between the Fourier modes in σ . Note that the $AdS_5 \times S^5$ superparticle is effectively the special case of this limit: while sending $T \rightarrow \infty$ implies that σ -derivatives are constrained to be zero, their contribution is dropped in the $T \rightarrow 0$ limit.

Solving for the momenta in the phase space action (5.23) one finds the light-cone Lagrangian corresponding to the Hamiltonian (5.24), which generalizes (4.18). It is useful to present it in terms of the “conformally-flat” 10-d coordinates (x^a, Z^M) , where Z^M is the “inverse” of Y^M . The corresponding form of the $AdS_5 \times S^5$ metric is

$$ds^2 = R^2 Z^{-2} (dx^a dx^a + dZ^M dZ^M), \quad Z^M = Y^{-2} Y^M = e^{-\phi} u^M. \quad (5.26)$$

The full superstring Lagrangian expressed in terms of these coordinates was given in [2] and in Appendix C in [3]. Its phase space counterpart has the same structure as (5.23), (5.24) [3]

$$\mathcal{L} = \mathcal{P}_\perp \dot{x}_\perp + \mathcal{P}_M \dot{Z}^M + \frac{i}{2} p^+ (\theta^i \dot{\theta}_i + \eta^i \dot{\eta}_i - h.c.) - \mathcal{H}, \quad (5.27)$$

$$\begin{aligned} \mathcal{H} = & \frac{1}{2p^+} [\mathcal{P}_\perp^2 + \mathcal{P}_M \mathcal{P}_M + T^2 Z^{-4} (\dot{x}_\perp^2 + \dot{Z}^M \dot{Z}^M) \\ & + Z^{-2} [p^{+2} (\eta^2)^2 + 2ip^+ \eta \rho^{MN} \eta Z_M \mathcal{P}_N]] \\ & - T [|Z|^{-3} \eta^i \rho_{ij}^M Z^M (\dot{\theta}^j - i\sqrt{2}|Z|^{-1} \eta^j \dot{x}) + h.c.], \end{aligned} \quad (5.28)$$

with $p^+ \dot{x}^- + \mathcal{P}_\perp \dot{x}_\perp + \mathcal{P}_M \dot{Z}^M + \frac{i}{2} p^+ (\theta^i \dot{\theta}_i + \eta^i \dot{\eta}_i - h.c.) = 0$. Eliminating the momenta $\mathcal{P}_\perp$ and $\mathcal{P}_M$ in (5.27) gives the following light-cone Lagrangian corresponding to the Hamiltonian density (5.28)

$$\begin{aligned} \mathcal{L} = & \frac{p^+}{2} \left[\dot{x}_\perp^2 + (\dot{Z}^M - i\eta_i \rho^{MNi} \eta^j Z_N Z^{-2})^2 + i(\theta^i \dot{\theta}_i + \eta^i \dot{\eta}_i - h.c.) \right. \\ & \left. - Z^{-2} (\eta^2)^2 - T^2 Z^{-4} (p^+)^{-2} (\dot{x}_\perp^2 + \dot{Z}^M \dot{Z}^M) \right] \\ & - T [|Z|^{-3} \eta^i \rho_{ij}^M Z^M (\dot{\theta}^j - i\sqrt{2}|Z|^{-1} \eta^j \dot{x}) + h.c.]. \end{aligned} \quad (5.29)$$

As in (4.18), the factors of p^+ can be absorbed into the rescaling $\sigma \rightarrow \sigma/p^+$ in the action. The $T \sim \sqrt{\lambda} \rightarrow 0$ limit of this Lagrangian is obtained by omitting the σ -derivative terms. Our final result is thus

$$\begin{aligned} \mathcal{L}_{\lambda \rightarrow 0} = & \frac{p^+}{2} \left[\dot{x}_\perp^2 + i(\theta^i \dot{\theta}_i + \theta_i \dot{\theta}^i) \right. \\ & \left. + (\dot{Z}^M - i\eta_i \rho^{MNi} \eta^j Z_N Z^{-2})^2 + i(\eta^i \dot{\eta}_i + \eta_i \dot{\eta}^i) - Z^{-2} (\eta^2)^2 \right]. \end{aligned} \quad (5.30)$$

This Lagrangian corresponds to the $T \rightarrow 0$ limit of the Hamiltonian (5.28) or (5.25). An interesting feature of (5.30) is that the two transverse 4-d coordinates $x_\perp$ and the 4 complex fermions θ_i corresponding to the linear part of the light-cone $\mathcal{N} = 4$ supersymmetry decouple from the rest of the variables.

As already mentioned above, (5.30) contains the action of the superparticle in $AdS_5 \times S^5$ as its part depending only on constant modes in σ . The important next problem is to

show that in addition to the type IIB $AdS_5 \times S^5$ supergravity multiplet, whose presence is implied by the correspondence with the superparticle action [5] and is expected since the masses of “protected” states do not depend on λ , the string spectrum following from quantization of (5.30) contains also massless higher spin particles in AdS_5 . Their light-cone description in AdS_d was given in [15, 16]: in the case of totally symmetric field ϕ_s one starts with totally symmetric traceless $SO(d-2)$ tensor $\phi_{I_1 \dots I_s}$ and decomposes it further into irreducible representations of $SO(d-3)$ as $\sum_{s'=0}^s \phi_{s'}$, $\phi_{s'} = (\phi_{i_1 \dots i_{s'}})$. Then, in the parametrization of AdS space as in (5.26) and after an appropriate rescaling, the fields $\phi_{s'}$ satisfy the following free equations $(\partial_x^2 + \partial_Z^2 - Z^{-2} A_{s'}) \phi_{s'} = 0$. Here $A_{s'} = (s' + \frac{d-5}{2})^2 - \frac{1}{4}$, i.e. $A_{s'} = s'^2 - \frac{1}{4}$ in the case of AdS_5 . These equations should appear as the mass shell conditions for the “lightest” string states corresponding to (5.25) and (5.30).^p

To conclude, we expect that the light-cone approach developed in [15, 5] and [2, 3] should be instrumental in proving the conjectured AdS/CFT correspondence [7, 8] in the $\lambda \rightarrow 0$ limit.

Acknowledgments

This work was supported in part by the DOE grant DE-FG02-91ER-40690, PPARC grant PPA/G/S/1998/00613, INTAS project 991590 and CRDF RPI-2108 grant. We are grateful to S. Frolov, I. Klebanov, A. Mikhailov, H. Nastase, W. Siegel, M. Vasiliev and especially R. Metsaev for useful discussions.

References

- [1] J. Maldacena, “The large-N limit of superconformal field theories and supergravity,” Adv. Theor. Math. Phys. **2**, 231 (1998), [arXiv:hep-th/9711200]. S.S. Gubser, I.R. Klebanov and A.M. Polyakov, “Gauge theory correlators from non-critical string theory,” Phys. Lett. B **428**, 105 (1998), [arXiv:hep-th/9802109]. E. Witten, “Anti-de Sitter space and holography,” Adv. Theor. Math. Phys. **2**, 253 (1998), [arXiv:hep-th/9802150].
- [2] R.R. Metsaev and A.A. Tseytlin, “Superstring action in $AdS_5 \times S^5$: κ -symmetry light cone gauge,” Phys. Rev. D **63**, 046002 (2001) [arXiv:hep-th/0007036].
- [3] R.R. Metsaev, C.B. Thorn and A.A. Tseytlin, “Light-cone Superstring in AdS Space-time,” Nucl. Phys. B **596**, 151 (2001) [arXiv:hep-th/0009171].
- [4] M.B. Green and J.H. Schwarz, “Supersymmetric String Theories,” Phys. Lett. B **109**, 444 (1982).
- [5] R.R. Metsaev, “Light cone gauge formulation of IIB supergravity in $AdS_5 \times S^5$ background and AdS/CFT correspondence,” Phys. Lett. B **468**, 65 (1999) [arXiv:hep-th/9908114].

^pThe non-normalizable and normalizable solutions for $\phi_s(x, z)$ which should represent the conformal higher spin fields and the associated conformal higher spin currents of the boundary theory have indeed the right integer dimensions [15]: $\Delta = 2 - s$ and $\Delta = s + d - 3 = s + 2$.

- [6] U. Lindstrom, B. Sundborg and G. Theodoridis, “The Zero tension limit of the superstring,” *Phys. Lett.* **B253**, 319 (1991). J. Isberg, U. Lindstrom and B. Sundborg, “Space-time symmetries of quantized tensionless strings,” *Phys. Lett.* **B293**, 321 (1992) [arXiv:hep-th/9207005]. J. Isberg, U. Lindstrom, B. Sundborg and G. Theodoridis, “Classical and quantized tensionless strings,” *Nucl. Phys.* **B411**, 122 (1994) [arXiv:hep-th/9307108].
- [7] B. Sundborg, “Stringy gravity, interacting tensionless strings and massless higher spins,” *Nucl. Phys. Proc. Suppl.* **102**, 113 (2001) [arXiv:hep-th/0103247]. P. Haggi-Mani and B. Sundborg, “Free large N supersymmetric Yang-Mills theory as a string theory,” *JHEP* **0004**, 031 (2000) [arXiv:hep-th/0002189].
- [8] E. Witten, talk at J.H. Schwarz 60-th Birthday Conference, CalTech, November 2-3, 2001.
- [9] M.A. Vasiliev, “Conformal higher spin symmetries of 4D massless supermultiplets and $osp(L,2M)$ invariant equations in generalized (super)space,” arXiv:hep-th/0106149. “Cubic interactions of bosonic higher spin gauge fields in $AdS(5)$,” *Nucl. Phys. B* **616**, 106 (2001) [arXiv:hep-th/0106200].
- [10] E. Sezgin and P. Sundell, “Towards massless higher spin extension of $D = 5$, $N = 8$ gauged supergravity,” *JHEP* **0109**, 025 (2001) [arXiv:hep-th/0107186]. “Doubletons and 5D higher spin gauge theory,” *JHEP* **0109**, 036 (2001) [arXiv:hep-th/0105001].
- [11] A. Mikhailov, “Notes on higher spin symmetries,” arXiv:hep-th/0201019.
- [12] S.E. Konstein, M.A. Vasiliev and V.N. Zaikin, “Conformal higher spin currents in any dimension and AdS/CFT correspondence,” *JHEP* **0012**, 018 (2000) [arXiv:hep-th/0010239]. D. Anselmi, “Theory of higher spin tensor currents and central charges,” *Nucl. Phys. B* **541**, 323 (1999) [arXiv:hep-th/9808004]. “Higher-spin current multiplets in operator-product expansions,” *Class. Quant. Grav.* **17**, 1383 (2000) [arXiv:hep-th/9906167].
- [13] H. Nastase and W. Siegel, “A new AdS/CFT correspondence,” *JHEP* **0010**, 040 (2000) [arXiv:hep-th/0010106].
- [14] H. Liu and A.A. Tseytlin, “ $D = 4$ super Yang-Mills, $D = 5$ gauged supergravity, and $D = 4$ conformal supergravity,” *Nucl. Phys. B* **533**, 88 (1998) [arXiv:hep-th/9804083].
- [15] R.R. Metsaev, “Light cone form of field dynamics in anti-de Sitter spacetime and AdS/CFT correspondence,” *Nucl. Phys. B* **563**, 295 (1999) [arXiv:hep-th/9906217].
- [16] R.R. Metsaev, “Light-cone form of field dynamics in AdS space-time,” arXiv:hep-th/0011112.
- [17] N. Boulanger, M. Henneaux and P. van Nieuwenhuizen, “Conformal (super)gravities with several gravitons,” arXiv:hep-th/0201023.
- [18] E.S. Fradkin and A.A. Tseytlin, “Conformal Supergravity,” *Phys. Rept.* **119**, 233 (1985).
- [19] M. Kaku, P.K. Townsend and P. van Nieuwenhuizen, “Properties Of Conformal Supergravity,” *Phys. Rev. D* **17**, 3179 (1978).

- [20] E.S. Fradkin and V.Y. Linetsky, “A Superconformal Theory Of Massless Higher Spin Fields In $D = (2+1)$,” *Mod. Phys. Lett. A* **4**, 731 (1989); *Annals Phys.* **198**, 293 (1990).
- [21] E.S. Fradkin and V.Y. Linetsky, “Cubic Interaction In Conformal Theory Of Integer Higher Spin Fields In Four-Dimensional Space-Time,” *Phys. Lett. B* **231**, 97 (1989). “Superconformal Higher Spin Theory In The Cubic Approximation,” *Nucl. Phys. B* **350**, 274 (1991).
- [22] E.S. Fradkin and A.A. Tseytlin, “Conformal Anomaly In Weyl Theory And Anomaly Free Superconformal Theories,” *Phys. Lett. B* **134**, 187 (1984).
- [23] R.R. Metsaev and A.A. Tseytlin, “Type IIB superstring action in $AdS_5 \times S^5$ background,” *Nucl. Phys. B* **533**, 109 (1998), hep-th/9805028.
- [24] R. Kallosh and A. Rajaraman, “Vacua of M-theory and string theory,” *Phys. Rev. D* **58**, 125003 (1998) [arXiv:hep-th/9805041].
- [25] I. Pesando, “A kappa gauge fixed type IIB superstring action on $AdS_5 \times S^5$,” *JHEP* **9811**, 002 (1998), [arXiv:hep-th/9808020]. “All roads lead to Rome: Supersolvable and supercosets,” *Mod. Phys. Lett. A* **14**, 343 (1999) [arXiv:hep-th/9808146].
- [26] R. Kallosh and J. Rahmfeld, “The GS string action on $AdS_5 \times S^5$,” *Phys. Lett. B* **443**, 143 (1998), [arXiv:hep-th/9808038]. R. Kallosh and A.A. Tseytlin, “Simplifying superstring action on $AdS_5 \times S^5$,” *JHEP* **10**, 016 (1998), [arXiv:hep-th/9808088].
- [27] R. Roiban and W. Siegel, “Superstrings on $AdS(5) \times S(5)$ supertwistor space,” *JHEP* **0011**, 024 (2000) [arXiv:hep-th/0010104].
- [28] A.M. Polyakov, “Gauge fields and space-time,” arXiv:hep-th/0110196.
- [29] A.M. Polyakov, private communication (Jan. 2000).
- [30] A.A. Tseytlin, “Superstrings in AdS in light cone gauge,” *Int. J. Mod. Phys. A* **16**, 900 (2001) [arXiv:hep-th/0009226].
- [31] J. Polchinski and L. Susskind, “String theory and the size of hadrons,” arXiv:hep-th/0112204.